

# TRINITY™

*Diascia barberae*



**TRINITY™ Grace**

- Large flowers and exceptional weather tolerance
- Holds blooms longer for an improved shelf life
- Grows into a small bush
- Mature Height and Width 12 inches X 14 inches (30cm x 36cm)



**TRINITY™ Sunset**

For additional technical growing information:

TRINITY™ [CLICK HERE](#)



 **DANZIGER**

[www.danzigeronline.com](http://www.danzigeronline.com) |   

# TRINITY™

Diascia barberae



## PROPAGATION TIPS

- Stick on priority – Number 3 out of 4 categories



Flowering time

Spring / Summer  
Flowering



Width

(12 - 14")  
30 - 36 cm



Height

(14 - 16")  
36 - 41 cm

Average Time	Temperature	Hormone	Fertilization	Fungicide
5 weeks	Weeks 1 - 2 70°-72° F (21° - 22°C)	Optional	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 5 68° - 70° (20° - 21° C)		Weeks 3-5 100 to 150 ppm N	Apply broad spectrum Week 2



**Misting specification** - Remove from mist as soon as possible  
- Spray adjuvant at sticking



- Maintain warm temperatures allowing plants to develop.
- Spray Daminozide at 1500 to 2000 ppm
- Pinch is recommended week 4 or at transplant

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.8 / 6.2	0.80	72 / 74 F (22 / 23 C)	50 ppm N in mist	Not required		Day of sticking	
Week 2	5.8 to 6.2	.80 to .90	72 / 74 F (22 / 23 C)	50 ppm N in mist	Not required		Second fungicide app	
Week 3	5.8 to 6.2	.90 to 1.0	72 / 74 F (22 / 23 C)	Feed 100 ppm to 150 ppm	Not required			
Week 4	5.8 to 6.2	.90 to 1.0	Cool to 68 / 70 F (20 / 21 C)	Feed 100 to 150 ppm	Not required	Daminozide as needed		Pinch
Week 5	5.8 to 6.2	0.9 to 1.0	68 F (20 C)	Feed 100 to 150 ppm	Not required	Daminozide as needed		

For additional technical growing information:

TRINITY™ [CLICK HERE](#)



**DANZIGER**

[www.danzigeronline.com](http://www.danzigeronline.com)



# TRINITY™

Diascia barberae



## FINISHING TIPS

- Pinch week 4 in liner or at transplant
- Plant liners above new soil – not to bury plant meristem

Average Time (from liners)	Temperature	Pinch/ Daylength Modification	Fertilization	Plant Growth Regulator
8 to 9 wks	Average Day	Pinch –Yes	100 - 150 ppm N	In liner and finish stages
(12 cm) 5” pot	65° - 68° (18° - 20° C)			
8 to 10 wks	65° - 68° (18° - 20° C)	Day Neutral	Soil EC 1.0 - 1.2 pH 5.8 to 6.2	
(15 cm) 6 ½”				

<b>Pests</b>	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Thrip	METHIOCARB, ACRINATHRIN, ABAMECTIN, DICHLORVOS, SPINOSAD
<b>Diseases</b>	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRADIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB



- PGR - Spray Daminozide at 2500 to 3750 ppm for finishing plants
- Paclobutrazol – Do no apply
- Spray broad spectrum fungicide after transplant
- Scout for Aphids and Thrip on regular basis

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
<b>Week 6 Transplant</b>	5.8 to 6.2	0.9 to 1.2	65 to 68 F (18 to 20 C)	feed 100 to 150 ppm		Daminozide as needed	Spray fungicide after transplant	
<b>Week 6 to 12</b>	5.8 to 6.2	0.9 to 1.2	65 to 68 F (18 to 20 C)	feed 100 to 150 ppm				Do not apply Paclobutrazol

For additional technical growing information:

TRINITY™ [CLICK HERE](#)



 **DANZIGER**

[www.danzigeronline.com](http://www.danzigeronline.com) |   